

# ANCHOR PRESENTATIONS AT 2019 AAOS ANNUAL MEETING

## TUESDAY-WEDNESDAY POSTER

### Poster No. P0043

#### Surgical Treatment of Femoroacetabular Impingement: Arthroscopy vs. Surgical Hip Dislocation – A Propensity Matched Analysis

*Jeffrey J. Nepple, MD, Saint Louis, MO*

*Ira Zaltz, MD, Royal Oak, MI*

*Asheesh Bedi, MD, Ann Arbor, MI*

*Paul E. Beaulé, MD, Ottawa, ON, Canada*

*Michael B. Millis, MD, Boston, MA*

*Rafael J. Sierra, MD, Rochester, MN*

*Ernest L. Sink, MD, New York, NY*

*John C. Clohisey, MD, Saint Louis, MO*

*ANCHOR Study Group*

In a propensity matched analysis, patients undergoing hip arthroscopy or surgical hip dislocation demonstrated similar outcomes.

## THURSDAY-SATURDAY POSTER

### Poster No. P0913

#### Predictors of Failure After Surgical Treatment of Femoroacetabular Impingement: Results of a Multicenter Prospective Cohort of 621 Hips

*Jeffrey J. Nepple, MD, Saint Louis, MO*

*Asheesh Bedi, MD, Ann Arbor, MI*

*Ira Zaltz, MD, Royal Oak, MI*

*Christopher M. Larson, MD, Edina, MN*

*Paul E. Beaulé, MD, Ottawa, ON, Canada*

*Ernest L. Sink, MD, New York, NY*

*Young Jo Kim, MD, PhD, Boston, MA*

*John C. Clohisey, MD, Saint Louis, MO*

*ANCHOR Study Group*

Several patient and disease characteristics were independently associated with failure and may help guide patient-specific outcome expectations of FAI surgery based on results of a multicenter study.

## WEDNESDAY EDUCATIONAL PROGRAM

**2:00 PM**

**PAPER 395**

### Gender Differences in Outcome after Corrective Surgery for Femoroacetabular Impingement (FAI) Reflect Differences in Preoperative Baseline Scores

*Tristan Maerz, PhD, Ann Arbor, MI*

*Geneva Baca, St. Louis, MO*

*Paul E. Beaulé, MD, Ottawa, ON, Canada*

*John C. Clohisey, MD, St. Louis, MO*

*Young Jo Kim, MD, PhD, Boston, MA*

*Christopher M. Larson, MD, Edina, MN*

*Michael B. Millis, MD, Boston, MA*

*David A. Podeszwa, MD, Dallas, TX*

*Perry L. Schoenecker, MD, St. Louis, MO*

*Rafael J. Sierra, MD, Rochester, MN*

*Ernest L. Sink, MD, New York, NY*

*Daniel J. Sucato, MD, MS, Dallas, TX*

*Robert T. Trousdale, MD, Rochester, MN*

*Ira Zaltz, MD, Royal Oak, MI*

*Asheesh Bedi, MD, Ann Arbor, MI*

*ANCHOR Multicenter Study Group*

In a large, multi-center, prospective cohort of FAI, gender differences can be traced to preoperative baseline scores.